

# UNCLASSIFIED

AD NUMBER
ADB274604
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies only; Proprietary Info.; May 2001. Other requests shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott St., Ft. Detrick, MD 21702-5012.
AUTHORITY
USAMRMC ltr, 26 Aug 2002

THIS PAGE IS UNCLASSIFIED

AD\_\_\_\_\_

Award Number: DAMD17-98-1-8149

TITLE: Molecular Analysis of the Cripto Growth Factor Receptor

PRINCIPAL INVESTIGATOR: Minjung Kim  
Chaosu E  
Michael M. Shen, Ph.D.

CONTRACTING ORGANIZATION: University of Medicine and Dentistry of New Jersey  
Robert Wood Johnson Medical School  
Piscataway, New Jersey 08854-5638

REPORT DATE: May 2001

TYPE OF REPORT: Final

PREPARED FOR: U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Distribution authorized to U.S. Government agencies only (proprietary information, May 2001). Other requests for this document shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott Street, Fort Detrick, Maryland 21702-5012.

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

20020124 388

## NOTICE

USING GOVERNMENT DRAWINGS, SPECIFICATIONS, OR OTHER DATA INCLUDED IN THIS DOCUMENT FOR ANY PURPOSE OTHER THAN GOVERNMENT PROCUREMENT DOES NOT IN ANY WAY OBLIGATE THE U.S. GOVERNMENT. THE FACT THAT THE GOVERNMENT FORMULATED OR SUPPLIED THE DRAWINGS, SPECIFICATIONS, OR OTHER DATA DOES NOT LICENSE THE HOLDER OR ANY OTHER PERSON OR CORPORATION; OR CONVEY ANY RIGHTS OR PERMISSION TO MANUFACTURE, USE, OR SELL ANY PATENTED INVENTION THAT MAY RELATE TO THEM.



### LIMITED RIGHTS LEGEND

Award Number: DAMD17-98-1-8149

Organization: University of Medicine and Dentistry of New Jersey  
Robert Wood Johnson Medical School

Those portions of the technical data contained in this report marked as limited rights data shall not, without the written permission of the above contractor, be (a) released or disclosed outside the government, (b) used by the Government for manufacture or, in the case of computer software documentation, for preparing the same or similar computer software, or (c) used by a party other than the Government, except that the Government may release or disclose technical data to persons outside the Government, or permit the use of technical data by such persons, if (i) such release, disclosure, or use is necessary for emergency repair or overhaul or (ii) is a release or disclosure of technical data (other than detailed manufacturing or process data) to, or use of such data by, a foreign government that is in the interest of the Government and is required for evaluational or informational purposes, provided in either case that such release, disclosure or use is made subject to a prohibition that the person to whom the data is released or disclosed may not further use, release or disclose such data, and the contractor or subcontractor or subcontractor asserting the restriction is notified of such release, disclosure or use. This legend, together with the indications of the portions of this data which are subject to such limitations, shall be included on any reproduction hereof which includes any part of the portions subject to such limitations.

THIS TECHNICAL REPORT HAS BEEN REVIEWED AND IS APPROVED FOR PUBLICATION.

  
\_\_\_\_\_  
\_\_\_\_\_

**REPORT DOCUMENTATION PAGE**Form Approved  
OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

<b>1. AGENCY USE ONLY (Leave blank)</b>		<b>2. REPORT DATE</b> May 2001	<b>3. REPORT TYPE AND DATES COVERED</b> Final (13 Apr 98 - 12 Apr 01)	
<b>4. TITLE AND SUBTITLE</b> Molecular Analysis of the Cripto Growth Factor Receptor			<b>5. FUNDING NUMBERS</b> DAMD17-98-1-8149	
<b>6. AUTHOR(S)</b>  Minjung Kim Chaosu E Michael M. Shen, Ph.D.				
<b>7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)</b>  University of Medicine and Dentistry of New Jersey Robert Wood Johnson Medical School Piscataway, New Jersey 08854-5638 E-Mail: <a href="mailto:mshen@cabm.rutgers.edu">mshen@cabm.rutgers.edu</a>			<b>8. PERFORMING ORGANIZATION REPORT NUMBER</b>	
<b>9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)</b>  U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012			<b>10. SPONSORING / MONITORING AGENCY REPORT NUMBER</b>	
<b>11. SUPPLEMENTARY NOTES</b> Report contains color graphics				
<b>12a. DISTRIBUTION / AVAILABILITY STATEMENT</b> Distribution authorized to U.S. Government agencies only (proprietary information, May 2001). Other requests for this document shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott Street, Fort Detrick, Maryland 21702-5012.				<b>12b. DISTRIBUTION CODE</b>
<b>13. ABSTRACT (Maximum 200 Words)</b> Over the past three years, I have been investigating the biochemical functions of the Cripto gene, which has been implicated in autocrine/paracrine signaling during human breast carcinogenesis. Since previous studies have suggested that Cripto encodes a growth factor-like molecule, I initially studied the binding of Cripto to mammary cell lines to examine the distribution of putative receptors. However, recent molecular genetic evidence has emerged that Cripto instead may act as a co-receptor for signaling by Nodal, a member of the transforming growth factor-beta family. To investigate this model at the biochemical level, I have expressed soluble forms of Cripto and Nodal in transfected mammalian cells, and have used co-immunoprecipitation approaches to demonstrate a direct binding interaction between Cripto and Nodal. These findings suggest that Cripto functions through modulation of Nodal signaling in mammary carcinogenesis.				
<b>14. SUBJECT TERMS</b> (1) growth factor receptor; (2) mammary gland development (3) histochemical affinity reagent; (4) ligand-receptor interaction				<b>15. NUMBER OF PAGES</b> 14
				<b>16. PRICE CODE</b>
<b>17. SECURITY CLASSIFICATION OF REPORT</b> Unclassified	<b>18. SECURITY CLASSIFICATION OF THIS PAGE</b> Unclassified	<b>19. SECURITY CLASSIFICATION OF ABSTRACT</b> Unclassified	<b>20. LIMITATION OF ABSTRACT</b> Unlimited	

NSN 7540-01-280-5500

Standard Form 298 (Rev. 2-89)  
Prescribed by ANSI Std. Z39-18  
298-102

## Table of Contents

Cover	
SF 298	Page 2
Table of Contents	Page 3
Introduction	Page 4
Body	Page 4
Key Research Accomplishments	Page 11
Reportable Outcomes	Page 11
Conclusions	Page 11
References	Page 11
Appendices	N/A

## Introduction:

Work in our laboratory on breast development and tumorigenesis has focused on the *Cripto* gene, which encodes an extracellular protein that is a member of the *EGF-CFC* gene family. Several lines of evidence have indicated that *Cripto* is involved in autocrine or paracrine signaling during human breast carcinogenesis (reviewed in (Salomon et al. 1999)). In particular, *Cripto* is overexpressed in a large percentage of human breast cancers, but not in normal breast tissue (Qi et al. 1994), and *Cripto* has transforming activity when overexpressed in NOG-8 mouse mammary epithelial cells (Ciardiello et al. 1991). To investigate the role of *Cripto* in breast development and tumorigenesis, I have been analyzing the molecular mechanisms of *Cripto* protein activity, with the objective of identifying a putative *Cripto* receptor(s).

## Body:

The function of *Cripto* was initially of interest because its overexpression was found in approximately 80% of human breast carcinomas, as well as in colorectal and pancreatic cancers. Previous studies have suggested that *Cripto* has growth factor activity, and that addition of exogenous *Cripto* protein to mammary epithelial cells lines can result in tyrosine phosphorylation of the ErbB4 receptor and activation of the Ras/MAPK signaling pathway. However, since *Cripto* does not bind to any of the four known members of the *ErbB* receptor family, it has been unclear how *Cripto* might function as a ligand to activate receptor-mediated signaling.

Over the past three years, I have made significant progress towards understanding the molecular mechanisms of *Cripto* function, which has contributed to a substantial revision of our initial view of *Cripto* encoding an EGF (epidermal growth factor)-related growth factor. Based on my work and recent data from others, it is now evident that *Cripto* may in fact encode a co-receptor rather than a ligand, and is essential for signaling by a divergent member of the TGF-beta (transforming growth factor-beta) superfamily known as Nodal.

## Review of recent findings in the field

Members of the *EGF-CFC* gene family include mammalian *Cripto* and *Cryptic*, frog *FRL-1*, and zebrafish *oep*, which encode proteins that share an N-terminal signal sequence, a variant EGF-like motif, a novel conserved cysteine-rich domain (the CFC (*Cripto*, *Frl-1*, and *Cryptic*) motif), and a C-terminal hydrophobic region (reviewed in (Shen and Schier 2000)). These genes encode extracellular proteins that are localized to the surface of transfected cells (Shen et al. 1997; Zhang et al. 1998). This association is mediated by the C-terminal hydrophobic domain, which in the case of *Cripto* is required for glycosyl-phosphatidylinositol (GPI) linkage to the cell membrane (Minichiotti et al. 2000). Notably, all *EGF-CFC* family members appear to have functionally similar activities in assays for phenotypic rescue of *oep* mutant fish embryos by mRNA microinjection (Gritsman et al. 1999).

Work from our laboratory has provided essential insights into the biological functions and potential biochemical activities of EGF-CFC proteins through analysis of knock-out mice for *Cripto* and *Cryptic* (reviewed in (Shen and Schier 2000)). We have found that *EGF-CFC* genes function in axis formation at early stages in the mouse embryo, so that *Cripto* is required for correct orientation of the A-P (anterior-posterior) axis, while *Cryptic* is necessary for determination of the L-R (left-right) axis (Ding et al. 1998; Yan et al. 1999). In contrast to previous models for *Cripto* having growth factor activity (Salomon et al. 1999), these and other recent lines of evidence indicate that EGF-CFC proteins act as essential co-factors for a signaling factor known as Nodal (reviewed in (Schier and Shen 2000)). The *Nodal* gene encodes a divergent member of the TGF-

beta superfamily, and displays a mutant phenotype similar to that for *Cripto* (Zhou et al. 1993; Conlon et al. 1994). The downstream signaling pathway for *Nodal* has primarily been suggested by our current understanding of TGF-beta signal transduction pathways (Massagué 1998), and by the similar and/or synergistic phenotypes for targeted disruption of *Nodal*, the type II activin receptor *ActRIIB*, the type I receptor *ActRIB*, and the cytoplasmic signal transducer *Smad2* (Schier and Shen 2000); however, there is no biochemical evidence at present that *Nodal* protein directly binds to and activates activin receptors.

Taken together, these findings support a model in which EGF-CFC proteins form membrane-associated components of a receptor complex, and mediate *Nodal* signaling through signal-transducing partner(s) that include activin receptors (Gritsman et al. 1999; Shen and Schier 2000). Based on the requirement of *EGF-CFC* activity for *Nodal* signaling, the cell-autonomy of *oep* function in zebrafish, and the localization of EGF-CFC proteins at the cell surface, EGF-CFC proteins may function as co-receptors for *Nodal*. Thus, EGF-CFC proteins might interact with *Nodal* proteins to form a complex that binds to Activin-like receptors; alternatively, EGF-CFC factors could modify or induce conformational changes in either *Nodal* signals and/or Activin-like receptors that allow them to interact. In principle, release of EGF-CFC proteins from the membrane might result in a freely diffusible protein, which could form part of a receptor complex on a neighboring cell that may not itself express the *EGF-CFC* gene, in effect acting as a signal. Furthermore, the downstream signaling effects documented for *Cripto* protein might represent cross-talk between EGF receptor and *Smad2* signaling pathways (de Caestecker et al. 1998; Kretzschmar et al. 1999).

#### Technical Objective I: Expression patterns of *Cripto* and *Cripto* receptor


**Strategy:** At the start of our work on the *Cripto* gene, the existing published data suggested that *Cripto* was a EGF-like growth factor that functions through a novel receptor. Therefore, to analyze the functions of *Cripto*, I originally proposed to survey the distribution of endogenous *Cripto* expression as well as *Cripto*-AP binding sites in normal mammary tissue obtained from virgin, pregnant, lactating, and involuting mice. Furthermore, it had been shown that human *Cripto* protein (h*Cripto*) can bind with high affinity to a range of mammary epithelial cell lines, including MDA-MB-453, SK-BR-3 human breast cancer cells and normal mouse HC-11 cells, and can activate p42/44 MAP kinase in these cell lines (Brandt et al. 1994; Kannan et al. 1997). These data suggested that these cell lines expressed putative *Cripto* receptors.

**Results:** To examine the expression of *Cripto* at various stages of mammary development, I employed RT-PCR and ribonuclease protection assays to analyze mammary glands from virgin, pregnant, lactating, and regressing stages. As described in my 1999 annual report, these results indicated that *Cripto* is expressed at extremely low levels throughout mammary development, with slightly elevated expression during pregnancy and lactation (data not shown). Consequently, future studies should employ a different approach to investigate the levels and distribution of *Cripto* transcripts in the mammary gland, using our gene targeted mice that contain a *LacZ* reporter gene "knocked-in" under the transcriptional control of the *Cripto* promoter (Ding et al. 1998). These mice will allow *Cripto* expression to be examined by beta-galactosidase staining of mammary glands of *Cripto-LacZ*<sup>+/-</sup> heterozygotes.

In order to study the interaction between *Cripto* protein and its binding sites on mammary epithelial cell lines as well as tissues, I generated a soluble receptor affinity reagent that consists of human *Cripto* protein fused to secreted human placental alkaline phosphatase (AP) (Fig. 3A). The AP-*Cripto* fusion protein can be effectively expressed and secreted into conditioned media of transiently transfected COS7 cells. The advantage of using a AP fusion protein as a receptor affinity

reagent is that it is non-radioactive and sensitive, and that the AP fusion protein containing conditioned media can be directly used to visualize receptor binding on cell lines by histochemical staining (Berger et al. 1988; Flanagan and Leder 1990). Since human placental AP is heat-stable, endogenous AP activity can be readily inactivated by heating.

Using these AP-Cripto conditioned media, I established quantitative assays for Cripto binding to putative receptors. In this assay, I demonstrated that AP-hCripto protein specifically binds to the human mammary cell lines MDA-MB-453 and SK-BR-3 and that the binding is saturable (Fig. 3B). Scatchard analysis using MDA-MB-453 cells reveals a  $K_d$  of 10 nM for AP-hCripto binding to its binding site and approximately  $2.4 \times 10^5$  binding sites present on each cell (Fig. 3C).




**Figure 3. Binding of AP-Cripto to cultured cells.** (A) Schematic map of *pAPtag-4* expression vector. The *Cripto* gene was fused in frame to the 3' end of a DNA sequence encoding the intact secretory placental alkaline phosphatase. The CMV promoter can direct high level expression of the fusion protein gene in COS7 cells. (B) Quantitative binding of AP-Cripto to human breast cancer cell lines. Binding assays were performed using MDA-MB-453 and SK-BR-3 cells and AP-hCripto conditioned media. AP conditioned media prepared using the *pAPtag-4* vector was used as a control. The AP activities of both AP-hCripto and AP conditioned media were adjusted to same level (750U/ml). Each assay was performed in triplicate. (C) Scatchard analysis of AP-hCripto binding to its receptor on MDA-MB-453 cells. Quantitative binding assays were performed using serial AP-hCripto conditioned media dilutions with different concentrations of AP activity. Each concentration was assayed in triplicate. Scatchard analysis indicated a  $K_d$  of 10 nM and  $2.4 \times 10^5$  receptors per cell. Calculations were based on 1 pmol AP fusion protein equal to 30 units AP activity (Cheng and Flanagan 1994). Primary data are shown in inset.

**Heparin binding of Cripto.** Since our laboratory focuses on mouse models, I was particularly interested in examining binding to the mouse Cripto receptor(s). For this purpose, mouse mammary HC-11 cells and AP fusions with mouse Cripto protein (AP-mCripto) were used for binding assays. Surprisingly, I found that while AP-hCripto can bind to HC-11 cells with high affinity, AP-mCripto did not (Fig. 4A). In particular, the level of AP-mCripto binding to HC-11 cells was at least 20-fold lower than that of AP-hCripto.

Because some growth factors in the EGF family have been shown to bind heparin, I investigated the possibility that human Cripto could bind to heparan sulfate, a common component of the extracellular matrix, while mouse Cripto did not. My results from a heparin bead binding assay showed that this was the case (Figure 4B). The AP-hCripto strongly binds to heparin beads, but the binding of AP-mCripto was similar to that for the AP4 control. Therefore, these data indicate that human Cripto and mouse Cripto differ in their ability to bind heparin.






**Figure 4. Quantitative binding assay. (A) Binding of mouse Cripto and human Cripto to HC-11 Cells.**  $3.5 \times 10^5$  HC-11 cells were suspended in 1 ml AP-mCripto or AP-hCripto conditioned media (2000 U/ml). After incubation and wash, the cells were lysed. The cell lysates were heated at  $65^\circ\text{C}$  for 10 minute to inactivate endogenous AP activity and then the bound AP activity was detected as described (Flanagan and Leder 1990). AP conditioned media (2000 U/ml) was used as a control. **(B) Binding of mouse Cripto and human Cripto to heparin-beads.** 0.15 ml heparin-beads were mixed with 1 ml AP-mCripto or AP-hCripto conditioned media (1000 U/ml) and incubated at room temperature for 60 minutes. After washing with 1xPBS, the bound AP-Cripto was eluted with 0.3 ml 1xPBS/0.35M NaCl. The AP activity in the eluates was assayed. AP4 conditioned media (1000 U/ml) was used as a control.

Based on these findings, I concluded that the binding sites detected on HC-11 cells for human Cripto are likely to correspond to heparan sulfate proteoglycans, and not to signal-transducing receptors. Therefore, I did not pursue further AP-Cripto binding studies on mammary tissue, as originally proposed, but instead focused on biochemical studies of Cripto binding interactions. In summary, my studies addressed Task 1 of Technical Objective I in the original Statement of Work, but the lack of evidence for authentic receptor interactions did not support further studies in several lines of MMTV transgenic mice (Task 2 of Technical Objective I in the Statement of Work). In the meantime, the findings from our lab and others that Cripto is likely to function as a co-receptor for Nodal led me to focus on analyzing biochemical interactions with Nodal, as described below.

### Technical Objective II: Cloning and characterization of the *Cripto* receptor

**Strategy:** In my original proposal and Statement of Work, I described a methodology to clone putative Cripto receptors using AP-fusion protein reagents to screen expression libraries from HC-11 and MD-MBA-453 cells. However, based on our heparin binding studies described above, as well as on preliminary expression library screening results that were negative (data not shown), I decided to pursue an alternative approach to identify Cripto binding proteins.

**Results:** In my initial experiments, I hypothesized that the failure to detect the binding of AP-mCripto to HC-11 cells might instead be due to the low sensitivity of the AP binding assay. Therefore, I decided to use AP-mCripto conditioned media and radio-labeled HC-11 cell lysate to perform a sensitive pull-down assay to identify mouse Cripto-specific binding proteins.



**Figure 5. AP pull-down assay to identify mouse Cripto binding protein.** (A) Pull-down assay using radio-labeled HC-11 cell lysate. 0.5 ml radio-labeled HC-11 lysate were mixed with 0.3 ml AP-mCripto conditioned media or AP4 conditioned media (2500 U/ml) as a control. After incubation at 4°C for 60 minutes, AP-mCripto or AP were immunoprecipitated with 5 µl AP monoclonal antibody and 30 µl Staph A cells. The precipitates were resolved by 10% SDS-PAGE. (B) Pull-down assay using biotin-labeled HC-11 cell lysate. HC-11 cells were labeled with 0.25 mg/ml sulfo-NHS-LC-biotin which only biotinylates cell surface proteins and then were lysed with 1xRIPA buffer. The immunoprecipitation assay was performed as for radio-labeled HC-11 cells. The immunoprecipitates were resolved by 10% SDS-PAGE, transferred onto PVDF membrane, and detected with avidin-HRP conjugate (Altin and Pagla 1995).

In this pull-down assay,  $^{35}\text{S}$ -methionine labeled HC-11 cell lysate was mixed with AP-mCripto conditioned media or AP conditioned media as a control, and incubated at 4°C for 1 hour. AP monoclonal antibody was used to immunoprecipitate AP-mCripto or AP4 proteins, and the radio-labeled proteins co-immunoprecipitated with AP-mCripto or AP4 were identified by running a SDS-PAGE and exposing to X-ray film. Two protein bands, a major 70 Kd band and a weaker and slightly lower band, were detected for specifically binding to AP-mCripto but not AP4 (Fig. 5A). This finding suggested that these two proteins might be mouse Cripto-specific binding proteins.


To ask whether the two proteins were cell surface proteins, I used sulfo-NHS-LC-biotin, which only biotinylates cell surface proteins, to label HC-11 cells (Altin and Pagla 1995). Using the biotin-labeled HC-11 cell lysates, the same pull-down assay as described for the radio-labeled lysate was performed. The immunoprecipitates were resolved by SDS-PAGE, transferred onto PVDF membrane, and detected with avidin-HRP conjugate. In this assay, two AP-mCripto binding proteins were identified with the same molecular weight as the two detected in the radio-labeled pull-down assay (Fig. 5B), suggesting they are cell surface proteins.

**Interaction of Nodal protein with Cripto and Oep proteins.** During the course of this work, genetic studies in our lab as well as others showed that EGF-CFC proteins are required for Nodal signaling. In particular, Cripto and Oep proteins have been shown to be membrane-associated proteins. Based on these findings, it was suggested that EGF-CFC proteins act as essential co-factors or co-receptors for Nodal signaling, perhaps by mediating the binding of Nodal to its signal transducing receptors (type I and II activin receptors) through protein-protein interactions. Therefore, I decided to investigate the potential interaction between Nodal and EGF-CFC proteins.

For this purpose, I made constructs for expressing C-terminal truncated CRIPTO (mCRA) containing HA or Flag epitope tags; the C-terminally truncated Cripto and Oep constructs were

used to obtain efficient secretion into conditioned media. My results showed that Cripto and Oep proteins can be effectively produced by 293T cells; however, the mCRA-HA proteins expressed from the pcDNA3 vector differs from that produced using pSecTagB vector, which contains an efficient signal sequence cleavage site. Thus, pcDNA3 expressed mCRA-HA showed a single 17 Kd major band on a Western blot, while pSecTagB expressed mCRA-HA had four bands ranging from 15-20 Kd in size (Fig. 6A), perhaps because the pSecTagB vector uses the heterologous Ig K chain signal sequence, which may result in different protein processing. I chose to use the pSecTagB vector for protein expression because it resulted in a significantly higher expression level than pcDNA3 (Fig. 6A). The Flag-Oep m134 protein produced by 293T cells migrated as a doublet with a 14-16 Kd sizes (Fig.6B).


In addition, I also showed that Nodal-Flag can be processed and secreted by 293T cells. The conditioned media produced by 293T cells expressing Nodal-Flag contained unprocessed 38 Kd Nodal as well as processed, mature 14 Kd Nodal proteins (Fig. 6C). In contrast, when I used COS7 cells to express Nodal protein, only unprocessed Nodal protein can be detected in cell lysates and not in the conditioned media, suggesting that Nodal protein processing and secreting is cell-line dependent (Fig. 6D). Furthermore, I also found that the Nodal-Flag protein was largely aggregated in the conditioned media of 293T cells, since most of the Nodal-Flag protein migrated at apparent molecular weights of over 200 Kd on non-reducing SDS-PAGE gels. However, other data from sucrose gradient analysis as well as biological activity assays in the chick suggested that there was Nodal activity present in these conditioned media, so I used these for further analysis in co-immunoprecipitation experiments.



**Figure 6. Expression of Nodal, Cripto, and Oep proteins by 293T or COS7 cells.** Transiently transfected 293T or COS7 cells were cultured in serum free Optimem medium, and protein expression levels in conditioned media were examined by Western blotting. *pcDNA3* transfected culture supernatant was used

as a control. (A) **Expression of mCRA-HA protein.** *pcDNA3/mCRA-HA* and *pSecTagB/mCRA-HA* were transfected into 293T cells and conditioned media collected. mCRA-HA protein were detected by Western blotting (10  $\mu$ l conditioned media/lane) with anti-HA monoclonal antibody (Invitrogen). (B) **Expression of Flag-Oep protein.** *pFlag-Oep<sup>m134</sup>* was used to express C-terminally-truncated Oep in 293T conditioned media, followed by detection by Western blotting (10  $\mu$ l conditioned media/lane) using anti-Flag monoclonal antibody (Sigma). (C) **Expression of Nodal-Flag in 293T cells.** *pcDNA3/Nodal-Flag* was used to express Nodal in 293T conditioned media; Nodal expression was detected by Western blotting (16  $\mu$ l/lane) using anti-Flag monoclonal antibody. (D) **Expression of Nodal-Flag by COS7 cells.** was transfected into COS7 cells, followed by Western blotting of conditioned media and cell lysates with anti-Flag monoclonal antibody (15  $\mu$ l conditioned media or cell lysate/lane). (E) **Aggregation of Nodal-Flag.** Conditioned media (8  $\mu$ l) from 293T cells transfected with *pcDNA3/Nodal-Flag* was analyzed on a non-reducing SDS-PAGE gel, followed by Western blotting with anti-Flag monoclonal antibody.

Consequently, I used conditioned media containing epitope-tagged Nodal, Cripto, and/or Oep proteins to perform co-immunoprecipitation assays to detect interaction of Nodal with Cripto or Oep. To detect interaction between Nodal and Cripto, mCRA-HA conditioned media expressed following transfected by *pcDNA3/mCRA-HA* or *pSecTagB/mCRA-HA* was mixed and incubated with Nodal-Flag conditioned media. mCRA-HA protein in the mixture was immunoprecipitated by adding anti-HA-beads, and immunoprecipitated complexes were resolved on a 13.5% SDS-PAGE gel and detected by Western blotting using anti-Flag monoclonal antibody. My results showed that Nodal-Flag can be co-immunoprecipitated with mCRA-HA (Fig. 7A), and similarly, Nodal-HA can be co-immunoprecipitated with Flag-Oep (Fig. 7B).



**Figure 7. Co-immunoprecipitation assays for detecting interaction between Nodal and EGF-CFC proteins.** (A) Interaction between Nodal and Cripto proteins. 0.32 ml mCRA-HA conditioned media was mixed with 0.64 ml Nodal-Flag conditioned media. After incubation, 20  $\mu$ l anti-HA monoclonal antibody (McAb) coupled beads were added to immunoprecipitate mCRA-HA protein. The immunoprecipitated complexes were resolved on a 13.5% SDS-PAGE gel, and Western blotting was performed using anti-Flag monoclonal antibody. (B) Interaction between Nodal and Oep proteins. 0.5 ml Flag-Oep<sup>m134</sup> and Nodal-HA conditioned media were co-immunoprecipitated using 60  $\mu$ l anti-Flag McAb-beads, followed by Western blotting using anti-HA monoclonal antibody.

In summary, based on my findings, I have provided biochemical evidence for a direct interaction between Nodal and EGF-CFC proteins. These experiments support the current model that Cripto represents a co-receptor for the Nodal ligand, and are not consistent with previous views of Cripto as a growth factor-like molecule. Thus, while I did not pursue the specific experiments in Tasks 1 and 2 of Technical Objective II in the original Statement of Work, I successfully

accomplished the identification of a Cripto-interacting protein (Task 3 of Technical Objective II in the Statement of Work), and thereby addressed the overall goal of this specific aim.

### Key Research Accomplishments:

- I have established an alkaline-phosphatase fusion protein system for assaying the interaction of Cripto with cell-surface proteins.
- I have demonstrated that mouse Cripto is a heparin-binding protein.
- I have expressed soluble Cripto and Nodal proteins in conditioned media from transfected mammalian cell lines for analyses of their activities and potential biochemical interactions.
- Co-immunoprecipitation analyses of these proteins has provided evidence of a direct binding interaction.
- My findings raise the possibility that *Cripto* overexpression in human breast cancer may result in de-regulation of TGF-beta signaling pathways.

### Reportable Outcomes:

#### *Published manuscripts:*

None.

#### *Meeting abstracts:*

Shen, M. M., E, C., Saplakoglu, U., Yan, Y.-T., and Ding, J. (2000). Functional analysis of *EGF-CFC* genes in mouse development indicates their essential role in Nodal signaling. Era of Hope Department of Defense Breast Cancer Research Program Meeting. (Atlanta, GA).

E, C., Saplakoglu, U., and Shen, M. M. (2000). Role for Cripto as a cofactor in the Nodal signaling pathway. Era of Hope Department of Defense Breast Cancer Research Program Meeting. (Atlanta, GA).

#### *Personnel supported by this grant:*

Chaosu E

Minjung Kim

#### *Degrees obtained supported by this award:*

Chaosu E, M.S. received 11/2000

### Conclusions:

Although *Cripto* was the first member of the *EGF-CFC* gene family to be identified, its biochemical function has been poorly understood. Thus, it has been significant that recent genetic data from mouse and zebrafish studies has indicated that EGF-CFC proteins function as co-factors

or co-receptors for the TGF-beta factor Nodal. In my work, I have expressed soluble Nodal and Cripto proteins in conditioned media from transfected mammalian cells to detect a physical interaction between Nodal and EGF-CFC proteins. My data support a direct association between Nodal and Cripto in vivo to facilitate Nodal signaling, presumably through activin receptors. At present, these data do not allow us to distinguish whether Cripto acts as a co-factor or as a co-receptor for Nodal, however our ongoing studies should be able to resolve this critical issue. Finally, this work suggests that the activity of soluble Cripto on mammary epithelial cell lines is due to activation of a Nodal signaling pathway, and indicate the necessity of further studies on Nodal signaling in breast carcinogenesis.

## References:

- Altin, J.G. and E.B. Pagla. 1995. A one-step procedure for biotinylation and chemical cross-linking of lymphocyte surface and intracellular membrane-associated molecules. *Anal. Biochem.* 244: 382-389.
- Berger, J., J. Hauber, R. Hauber, R. Geiger, and B.R. Cullen. 1988. Secreted placental alkaline phosphatase: a powerful new quantitative indicator of gene expression in eukaryotic cells. *Gene* 66: 1-10.
- Brandt, R., N. Normanno, W.J. Gullick, J.H. Lin, R. Harkins, D. Schneider, B.W. Jones, F. Ciardiello, M.G. Persico, F. Armenante, N. Kim, and D.S. Salomon. 1994. Identification and biological characterization of an epidermal growth factor-related protein: cripto-1. *J. Biol. Chem.* 269: 17320-17328.
- Cheng, H.-J. and J.G. Flanagan. 1994. Identification and cloning of ELF-1, a developmentally expressed ligand for the Mek4 and Sek receptor tyrosine kinases. *Cell* 79: 157-168.
- Ciardiello, F., R. Dono, N. Kim, M.G. Persico, and D.S. Salomon. 1991. Expression of cripto, a novel gene of the epidermal growth factor gene family, leads to in vitro transformation of a normal mouse mammary epithelial cell line. *Cancer Res.* 51: 1051-1054.
- Conlon, F.L., K.M. Lyons, N. Takaesu, K.S. Barth, A. Kispert, B. Herrmann, and E.J. Robertson. 1994. A primary requirement for nodal in the formation and maintenance of the primitive streak in the mouse. *Development* 120: 1919-1928.
- de Caestecker, M.P., W.T. Parks, C.J. Frank, P. Castagnino, D.P. Bottaro, A.B. Roberts, and R.J. Lechleider. 1998. Smad2 transduces common signals from receptor serine-threonine and tyrosine kinases. *Genes Dev.* 12: 1587-1592.
- Ding, J., L. Yang, Y.T. Yan, A. Chen, N. Desai, A. Wynshaw-Boris, and M.M. Shen. 1998. Cripto is required for correct orientation of the anterior-posterior axis in the mouse embryo. *Nature* 395: 702-707.
- Flanagan, J.G. and P. Leder. 1990. The kit ligand: a cell surface molecule altered in steel mutant fibroblasts. *Cell* 63: 185-194.
- Gritsman, K., J. Zhang, S. Cheng, E. Heckscher, W.S. Talbot, and A.F. Schier. 1999. The EGF-CFC protein one-eyed pinhead is essential for nodal signaling. *Cell* 97: 121-132.
- Kannan, S., M. De Santis, M. Lohmeyer, D.J. Riese II, G.H. Smith, N. Hynes, M. Seno, R. Brandt, C. Bianco, G. Persico, N. Kenney, N. Normanno, I. Martinez-Lacaci, F. Ciardiello, D.F. Stern, W.J. Gullick, and D.S. Salomon. 1997. Cripto enhances the tyrosine

- phosphorylation of Shc and activates mitogen-activated protein kinase (MAPK) in mammary epithelial cells. *J. Biol. Chem.* 272: 3330-3335.
- Kretschmar, M., J. Doody, I. Timokhina, and J. Massague. 1999. A mechanism of repression of TGFbeta/ Smad signaling by oncogenic Ras. *Genes Dev.* 13: 804-816.
- Massagué, J. 1998. TGF- $\beta$  signal transduction. *Ann. Rev. Biochem* 67: 753-791.
- Minchiotti, G., S. Parisi, G. Liguori, M. Signore, G. Lania, E.D. Adamson, C.T. Lago, and M.G. Persico. 2000. Membrane-anchorage of Cripto protein by glycosylphosphatidylinositol and its distribution during early mouse development. *Mech. Dev.* 90: 133-142.
- Qi, C.F., D.S. Liscia, N. Normanno, G. Merlino, G.R. Johnson, W.J. Gullick, F. Ciardiello, T. Sacki, R. Brandt, N. Kim, N. Kenney, and D.S. Salomon. 1994. Expression of transforming growth factor alpha, amphiregulin and cripto-1 in human breast carcinomas. *Br. J. Cancer* 69: 903-910.
- Salomon, D.S., C. Bianco, and M. De Santis. 1999. Cripto: a novel epidermal growth factor (EGF)-related peptide in mammary gland development and neoplasia. *Bioessays* 21: 61-70.
- Schier, A.F. and M.M. Shen. 2000. Nodal signalling in vertebrate development. *Nature* 403: 385-389.
- Shen, M.M. and A.F. Schier. 2000. The EGF-CFC gene family in vertebrate development. *Trends Genet.* 16: 303-309.
- Shen, M.M., H. Wang, and P. Leder. 1997. A differential display strategy identifies Cryptic, a novel EGF-related gene expressed in the axial and lateral mesoderm during mouse gastrulation. *Development* 124: 429-442.
- Yan, Y.-T., K. Gritsman, J. Ding, R.D. Burdine, J.D. Corrales, S.M. Price, W.S. Talbot, A.F. Schier, and M.M. Shen. 1999. Conserved requirement for EGF-CFC genes in vertebrate left-right axis formation. *Genes Dev.* 13: 2527-2537.
- Zhang, J., W.S. Talbot, and A.F. Schier. 1998. Positional cloning identifies zebrafish one-eyed pinhead as a permissive EGF-related ligand required during gastrulation. *Cell* 92: 241-251.
- Zhou, X., H. Sasaki, L. Lowe, B.L. Hogan, and M.R. Kuehn. 1993. Nodal is a novel TGF-beta-like gene expressed in the mouse node during gastrulation. *Nature* 361: 543-547.



DEPARTMENT OF THE ARMY

US ARMY MEDICAL RESEARCH AND MATERIEL COMMAND  
504 SCOTT STREET  
FORT DETRICK, MARYLAND 21702-5012

REPLY TO  
ATTENTION OF:

MCMR-RMI-S (70-1y)

26 Aug 02

MEMORANDUM FOR Administrator, Defense Technical Information  
Center (DTIC-OCA), 8725 John J. Kingman Road, Fort Belvoir,  
VA 22060-6218


SUBJECT: Request Change in Distribution Statement

1. The U.S. Army Medical Research and Materiel Command has reexamined the need for the limitation assigned to technical reports written for this Command. Request the limited distribution statement for the enclosed accession numbers be changed to "Approved for public release; distribution unlimited." These reports should be released to the National Technical Information Service.

2. Point of contact for this request is Ms. Kristin Morrow at DSN 343-7327 or by e-mail at Kristin.Morrow@det.amedd.army.mil.

FOR THE COMMANDER:

Encl

  
PHYLLIS M. RINEHART  
Deputy Chief of Staff for  
Information Management



ADB274369  
ADB256383  
ADB264003  
ADB274462  
ADB266221  
ADB274470  
ADB266221  
ADB274464  
ADB259044  
ADB258808  
ADB266026  
ADB274658  
ADB258831  
ADB266077  
ADB274348  
ADB274273  
ADB258193  
ADB274516  
ADB259018  
ADB231912  
ADB244626  
ADB256677  
ADB229447  
ADB240218  
ADB258619  
ADB259398  
ADB275140  
ADB240473  
ADB254579  
ADB277040  
ADB249647  
ADB275184  
ADB259035  
ADB244774  
ADB258195  
ADB244675  
ADB257208  
ADB267108  
ADB244889  
ADB257384  
ADB270660  
ADB274493  
ADB261527  
ADB274286  
ADB274269  
ADB274592  
ADB274604

ADB274596  
ADB258952  
ADB265976  
ADB274350  
ADB274346  
ADB257408  
ADB274474  
ADB260285  
ADB274568  
ADB266076  
ADB274441  
ADB253499  
ADB274406  
ADB262090  
ADB261103  
ADB274372